

Anthemisole ^{wsp}

Levamisole Hydrochloride 30% BP

Broad spectrum anthelmintic against round worms of GIT & Respiratory tract

Composition: Each gram powder contains-

Levamisole Hydrochloride BP 300 mg, Sucrose BP 690 mg

Pharmacology: Anthemisole acts by interfering carbohydrate metabolism of nematodes by inhibiting the production of succinate dehydrogenase. Inhibition of this enzyme causes complete and irreversible muscular paralysis of worm.

Indication: Anthemisole powder is used for the eradication of nematodes of alimentary tract e.g., *Ascaridia galli*, *Heterakis gallinarum*, *Capillaria* spp. and respiratory tract (e.g., *Syngamus trachea*). Anthemisole powder can also be used orally with water to eliminate fowl eye worm (e.g., *Oxyspirura mansoni*).

Dosage and Administration: 0.08g Anthemisole powder per kg body weight.

Or 1 day use according to the following dosage schedule

Poultry	Number	Dosage
Chick (4-6 weeks)	20	1g powder per L drinking water
Pullet (10 weeks)	10	
Laying Bird (Layer)	7	

Contraindication: The drug is contra-Indicated in animals hypersensitive to active ingredient.

Precaution: Once opened the sachet should be used quickly.

Side Effect: In therapeutic doses, there is no undesirable side effect of Anthemisole powder in chickens. Even egg production; fertility and hatchability are not adversely affected.

Storage: Store in a cool & dry place, protected from the light. Keep all medicine out of reach of children.

Presentation: Each box contains 100g×10 in alu foil sachet.

For Veterinary Use Only.